

Multicriteria Optimization

Some continuous and discrete dynamics

Guillaume Garrigos

Institut de Mathématiques et de Modélisation de Montpellier
Universidad Tecnica Federico Santa Maria

Sestri-Levante: Franco/Italian workshop
8-12 September 2014

- H is an Hilbert space,
- $f_i : H \rightarrow \mathbb{R}$ are Lipschitz continuous on bounded sets.
- $K \subset H$ is a closed convex non empty set of constraints,
- One of the objective functions is bounded from below.

- H is an Hilbert space,
- $f_i : H \rightarrow \mathbb{R}$ are Lipschitz continuous on bounded sets.
- $K \subset H$ is a closed convex non empty set of constraints,
- One of the objective functions is bounded from below.

One approach, the scalarization method :

chose $0 \leq \theta_i \leq 1$, $\sum_{i=1}^q \theta_i = 1$, and minimize $\sum_{i=1}^q \theta_i f_i$.

- H is an Hilbert space,
- $f_i : H \rightarrow \mathbb{R}$ are Lipschitz continuous on bounded sets.
- $K \subset H$ is a closed convex non empty set of constraints,
- One of the objective functions is bounded from below.

One approach, the scalarization method :

chose $0 \leq \theta_i \leq 1$, $\sum_{i=1}^q \theta_i = 1$, and minimize $\sum_{i=1}^q \theta_i f_i$.

We are looking for the **simultaneous** minimization of the f_i 's.

- 1 Multicriteria analysis
- 2 Continuous steepest descent dynamic

- 1 Multicriteria analysis
- 2 Continuous steepest descent dynamic

Directional derivative (of Clarke)

$$df(x; d) := \limsup_{\substack{t \downarrow 0 \\ x' \rightarrow x}} \frac{f(x' + td) - f(x')}{t}.$$

Subdifferential (of Clarke)

$$\partial f(x) := \{p \in H \mid \langle p, d \rangle \leq df(x; d) \forall d \in H\}.$$

Remark


If f is of class C^1 , then

$$\partial f(x) = \{\nabla f(x)\} \text{ and } df(x; d) = \langle \nabla f(x), d \rangle.$$

Tangent and normal cones

$$T_K(x) := \text{cl} \{d \in H \mid \exists \varepsilon > 0, \forall t \in]0, \varepsilon[, x + td \in K\}.$$

$$N_K(x) := \{p \in H \mid \langle p, d \rangle \leq 0 \forall d \in T_K(x)\}.$$




Descent direction


We say that $d \in H$ is a *descent direction* at x if $df_i(x; d) < 0$ holds for all $i = 1..q$.

We say that it is an *admissible* descent direction if moreover $d \in T_K(x)$.

Example



Example



Armijo direction

We say that a descent direction $d \in H$ is an *Armijo direction* if $\exists \varepsilon > 0$ s.t. for all $t \in]0, \varepsilon[$:

$$\forall i, f_i(x + td) < f_i(x) + \frac{t}{2} df_i(x; d).$$

We say that it is an *admissible* Armijo direction if moreover $x + td \in K$.

Pareto equilibrium(s)


Pareto equilibrium(s)

- We say that $x \in K$ is a Pareto if there is no $y \in K$ such that $\forall i f_i(y) \leq f_i(x)$ and $\exists l f_l(y) < f_l(x)$.

Pareto equilibrium(s)

- We say that $x \in K$ is a Pareto if there is no $y \in K$ such that $\forall i f_i(y) \leq f_i(x)$ and $\exists l f_l(y) < f_l(x)$.
- We say that $x \in K$ is a weak Pareto if there is no $y \in K$ s.t. $\forall i f_i(y) < f_i(x)$.


Example



Pareto equilibrium(s)

- We say that $x \in K$ is a Pareto if there is no $y \in K$ such that $\forall i f_i(y) \leq f_i(x)$ and $\exists l f_l(y) < f_l(x)$.
- We say that $x \in K$ is a weak Pareto if there is no $y \in K$ s.t. $\forall i f_i(y) < f_i(x)$.
- We say that $x \in K$ is a critical Pareto if $0 \in N_K(x) + \text{Conv}\{\partial f_i(x)\}$.

Example



Pareto equilibrium(s)

- We say that $x \in K$ is a Pareto if there is no $y \in K$ such that $\forall i f_i(y) \leq f_i(x)$ and $\exists l f_l(y) < f_l(x)$.
- We say that $x \in K$ is a weak Pareto if there is no $y \in K$ s.t. $\forall i f_i(y) < f_i(x)$.
- We say that $x \in K$ is a critical Pareto if $0 \in N_K(x) + \text{Conv}\{\partial f_i(x)\}$.

Pareto equilibrium(s)

- We say that $x \in K$ is a Pareto if there is no $y \in K$ such that $\forall i f_i(y) \leq f_i(x)$ and $\exists l f_l(y) < f_l(x)$.
- We say that $x \in K$ is a weak Pareto if there is no $y \in K$ s.t. $\forall i f_i(y) < f_i(x)$.
- We say that $x \in K$ is a critical Pareto if $0 \in N_K(x) + \text{Conv}\{\partial f_i(x)\}$.

Properties

- Pareto \Rightarrow weak Pareto \Rightarrow critical Pareto.
- If the f_i are convex, then weak Pareto \Leftrightarrow critical Pareto.
- If the f_i are strictly convex, then the 3 notions both coincide.

Proposition

The following statements are equivalent :

- x is a critical Pareto point,
- There is no admissible descent direction at x ,
- There is no admissible Armijo direction at x .

We will consider

- 1 a continuous dynamic $\dot{u}(t) = s(u(t))$, where $s : K \rightarrow H$ verify
 - $s(u) = 0$ if u is a critical Pareto point,
 - $s(u)$ is an admissible descent direction else.
- 2 an algorithm $u_{n+1} = u_n + t_n d_n$ where d_n is an admissible Armijo direction.


- 1 Multicriteria analysis
- 2 Continuous steepest descent dynamic

Definition

Given $x \in K$, the *multiobjective steepest descent direction* is

$$s(x) := - (N_K(x) + \text{Conv}\{\partial f_i(x)\})^0 .$$

Example



Definition

Given $x \in K$, the *multiobjective steepest descent direction* is

$$s(x) := - (N_K(x) + \text{Conv}\{\partial f_i(x)\})^0.$$

Definition

Given $x \in K$, the *multiobjective steepest descent direction* is

$$s(x) := - (N_K(x) + \text{Conv}\{\partial f_i(x)\})^0.$$

Obviously, x is a Pareto critical iff $s(x) = 0$.

The multiobjective steepest descent direction

Definition

Given $x \in K$, the *multiobjective steepest descent direction* is

$$s(x) := - (N_K(x) + \text{Conv}\{\partial f_i(x)\})^0.$$

Obviously, x is a Pareto critical iff $s(x) = 0$.

In a sense, $s(x)$ selects itself a different convex combination of the functions at each x .

The multiobjective steepest descent direction

Definition

Given $x \in K$, the *multiobjective steepest descent direction* is

$$s(x) := - (N_K(x) + \text{Conv}\{\partial f_i(x)\})^0.$$

Obviously, x is a Pareto critical iff $s(x) = 0$.

In a sense, $s(x)$ selects itself a different convex combination of the functions at each x .

Example

If $q = 1$, then $s(x) = \text{proj}_{T_K(x)}(-\nabla f(x))$.

The multiobjective steepest descent direction

Definition

Given $x \in K$, the *multiobjective steepest descent direction* is

$$s(x) := - (N_K(x) + \text{Conv}\{\partial f_i(x)\})^0.$$

Obviously, x is a Pareto critical iff $s(x) = 0$.

In a sense, $s(x)$ selects itself a different convex combination of the functions at each x .


Example

If $q = 1$, then $s(x) = \text{proj}_{T_K(x)}(-\nabla f(x))$.

Property

$s(x)$ is an admissible descent direction at x , whenever $s(x) \neq 0$.

Example



Why $s(x)$ is called the **steepest** descent ?

Why $s(x)$ is called the **steepest** descent ?

Recall that (one objective function, no constraint) :

$$\frac{-\nabla f(x)}{\|\nabla f(x)\|} = \operatorname{argmin}_{\|d\| \leq 1} df(x, d).$$

Why $s(x)$ is called the **steepest** descent ?

Recall that (one objective function, no constraint) :

$$\frac{-\nabla f(x)}{\|\nabla f(x)\|} = \operatorname{argmin}_{\|d\| \leq 1} df(x, d).$$

The multiobjective steepest descent direction generalizes this fact :

Theorem (Attouch, Garrigos, Goudou, 2014)

$$\frac{s(x)}{\|s(x)\|} = \operatorname{argmin}_{\|d\| \leq 1, d \in T_K(x)} \max_i df_i(x, d).$$


The Multi-Objective Gradient dynamic :

$$\text{(MOG) } \dot{u}(t) = s(u(t)) \text{ i.e } \dot{u}(t) + (N_K(u(t)) + \text{Conv}\{\partial f_i(u(t))\})^0 = 0$$

A continuous dynamic : example 1

(MOG) $\dot{u}(t) = s(u(t))$ i.e $\dot{u}(t) + (N_K(u(t)) + \text{Conv}\{\partial f_i(u(t))\})^0 = 0$


$$f_1(x) = \|x - a\|^2 \text{ and } f_2(x) = \|x - b\|^2$$



A continuous dynamic : example 2

$$\text{(MOG)} \quad \dot{u}(t) = s(u(t)) \text{ i.e. } \dot{u}(t) + (N_K(u(t)) + \text{Conv}\{\partial f_i(u(t))\})^0 = 0$$

$$f_1(x) = \frac{1}{2}\|x\|^2 \text{ and } f_2(x) = \langle a, x \rangle$$



$$\text{(MOG)} \quad \dot{u}(t) = s(u(t)) \text{ i.e. } \dot{u}(t) + (N_K(u(t)) + \text{Conv}\{\partial f_i(u(t))\})^0 = 0$$

Theorem (Attouch, Garrigos, Goudou, 2014)


Suppose that H is finite-dimensional, and that the functions are convex and bounded from below. Then for any $u_0 \in K$, there exists a strong solution $u : [0, +\infty[\rightarrow K$ of (MOG), such that $u(0) = u_0$.

Strong solution essentially means an absolutely continuous trajectory u satisfying (MOG) for a.e. $t > 0$.

A continuous dynamic : example 2

$$\text{(MOG)} \quad \dot{u}(t) = s(u(t)) \text{ i.e. } \dot{u}(t) + (N_K(u(t)) + \text{Conv}\{\partial f_i(u(t))\})^0 = 0$$

$$f_1(x) = \frac{1}{2}\|x\|^2 \text{ and } f_2(x) = \langle a, x \rangle$$



$$(MOG) \dot{u}(t) = s(u(t)) \text{ i.e. } \dot{u}(t) + (N_K(u(t)) + \text{Conv}\{\partial f_i(u(t))\})^0 = 0$$

Theorem (Attouch, Garrigos, Goudou, 2014)

Suppose that H is finite-dimensional, and that the functions are convex and bounded from below. Then for any $u_0 \in K$, there exists a strong solution $u : [0, +\infty[\rightarrow K$ of (MOG), such that $u(0) = u_0$.

The proof cannot rely on Cauchy-Lipschitz because of lack of Lipschitz regularity.

→ Use Morau-Yoshida's regularization onto the f_i 's and the indicator function.

→ Use Peano's existence theorem on the regularized system : it asks only continuity but do not guarantee uniqueness.

→ Pass to the limit. Hard.

The problem of uniqueness is still open.

Can we find hypotheses ensuring Lipschitz continuity of $s(u)$?

Local Lipschitz property

Suppose $K = H$, and that the functions are of class $C^{1,1}$.


The vector field s is Lipschitz continuous at u if :

- $q = 2$, and $\nabla f_1(u) \neq \nabla f_2(u)$.
- The vectors $\nabla f_i(u)$ are linearly independent.

A continuous dynamic : example 2

$$\text{(MOG)} \quad \dot{u}(t) = s(u(t)) \text{ i.e. } \dot{u}(t) + (N_K(u(t)) + \text{Conv}\{\partial f_i(u(t))\})^0 = 0$$

$$f_1(x) = \frac{1}{2}\|x\|^2 \text{ and } f_2(x) = \langle a, x \rangle$$



A continuous dynamic : Qualitative behaviour

Theorem (Attouch, Garrigos, Goudou, 2014)

Suppose that the objective functions are lower regular (convex, or continuously differentiable ...). Then for all $i = 1..q$, the function $t \mapsto f_i(u(t))$ is decreasing.

Theorem (Attouch, Garrigos, Goudou, 2014)

Suppose that the objective functions are lower regular (convex, or continuously differentiable ...). Then for all $i = 1..q$, the function $t \mapsto f_i(u(t))$ is decreasing.

Theorem (Attouch, Garrigos, Goudou, 2014)

Suppose that the objective functions are quasi-convex.

- Then any bounded trajectory is weakly convergent.
- The limit point is a weak Pareto if the functions are convex.
- The limit point is a critical Pareto if the functions are C^1 or convex, and under compact assumption on u .

Theorem (Attouch, Garrigos, Goudou, 2014)

Suppose that the objective functions are lower regular (convex, or continuously differentiable ...). Then for all $i = 1..q$, the function $t \mapsto f_i(u(t))$ is decreasing.

Theorem (Attouch, Garrigos, Goudou, 2014)

Suppose that the objective functions are quasi-convex.

- Then any bounded trajectory is weakly convergent.
 - The limit point is a weak Pareto if the functions are convex.
 - The limit point is a critical Pareto if the functions are C^1 or convex, and under compact assumption on u .
-
- We recover classic results by taking $q = 1$.

Theorem (Attouch, Garrigos, Goudou, 2014)

Suppose that the objective functions are lower regular (convex, or continuously differentiable ...). Then for all $i = 1..q$, the function $t \mapsto f_i(u(t))$ is decreasing.

Theorem (Attouch, Garrigos, Goudou, 2014)

Suppose that the objective functions are quasi-convex.

- Then any bounded trajectory is weakly convergent.
 - The limit point is a weak Pareto if the functions are convex.
 - The limit point is a critical Pareto if the functions are C^1 or convex, and under compact assumption on u .
-
- We recover classic results by taking $q = 1$.
 - Can we have strong convergence under stronger assumptions?


- A descent method associated to some scalarization $\sum_{i=1}^q \theta_i f_i$. In (MOG) the θ_i are chosen and modified automatically along the time. And ALL the functions decrease.

- A descent method associated to some scalarization $\sum_{i=1}^q \theta_i f_i$. In (MOG) the θ_i are chosen and modified automatically along the time. And ALL the functions decrease.
- A descent method associated to $\max f_j$.

A continuous dynamic : example 1

(MOG) $\dot{u}(t) = s(u(t))$ i.e $\dot{u}(t) + (N_K(u(t)) + \text{Conv}\{\partial f_i(u(t))\})^0 = 0$

$$f_1(x) = \|x - a\|^2 \text{ and } f_2(x) = \|x - b\|^2$$



The Multi-Objective Gradient dynamic

A Continuous Gradient-like Dynamical Approach to Pareto-Optimization in Hilbert Spaces. Attouch, Goudou, 2014

A Dynamic Gradient Approach to Pareto Optimization with Nonsmooth (...). Attouch, Garrigos, Goudou, Submitted.

Multi-Objective Gradient algorithm

Steepest descent methods for multicriteria optimization. Fliege, Svaiter, 2000.

A steepest descent method for vector optimization. Drummond, Svaiter, 2005.

Newton's method

Newton's Method for Multiobjective Optimization. Fliege, Drummond, Svaiter, 2009.

A quadratically convergent Newton method for vector optimization. Drummond, Raupp, Svaiter, 2014.

Quasi-Newton's method for multiobjective optimization. Povalej, 2014.

Proximal method

Proximal Methods in Vector Optimization. Bonnel, Iusem, Svaiter, 2005.

The Multi-Objective Gradient dynamic

A Continuous Gradient-like Dynamical Approach to Pareto-Optimization in Hilbert Spaces. Attouch, Goudou, 2014

A Dynamic Gradient Approach to Pareto Optimization with Nonsmooth (...). Attouch, Garrigos, Goudou, Submitted.

Multi-Objective Gradient algorithm

Steepest descent methods for multicriteria optimization. Fliege, Svaiter, 2000.

A steepest descent method for vector optimization. Drummond, Svaiter, 2005.

Newton's method

Newton's Method for Multiobjective Optimization. Fliege, Drummond, Svaiter, 2009.

A quadratically convergent Newton method for vector optimization. Drummond, Raupp, Svaiter, 2014.

Quasi-Newton's method for multiobjective optimization. Povalej, 2014.

Proximal method

Proximal Methods in Vector Optimization. Bonnel, Iusem, Svaiter, 2005.

Thank you for your attention !